



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

147. Proposed by R. A. WELLS, Professor of Mathematics, Franklin College, Athens, Ohio.

Find the locus in space of the point which is equally illuminated by each of two unequal lights whose intensities are a and b ($a > b$), placed at a distance c from each other.

Solution by WILLIAM HOOVER, A.M., Ph.D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Let (x, y, z) be the rectangular coördinates of any one of the illuminated points, the mid-point of the line joining the two lights taken along the x -axis as origin; then the coördinates of the two lights may be given as $(l, 0, 0)$, $(-l, 0, 0)$, and by theory

$$\frac{a}{(x-l)^2 + y^2 + z^2} = \frac{b}{(x+l)^2 + y^2 + z^2} \dots (1),$$

$$\text{or } x^2 + y^2 + z^2 + 2l \frac{a+b}{a-b} x + l^2 = 0 \dots (2), \text{ a sphere.}$$

Also solved by G. B. M. ZERR, and J. SCHEFFER.

148. Proposed by DR. E. D. ROE, JR., Associate Professor of Mathematics in Syracuse University, Syracuse, N. Y.

The condition that two triangles, abc, xyz , are similar is

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ x & y & z \end{vmatrix} = 0,$$

and the condition that the triangle abc is equilateral is

$$\begin{vmatrix} a & b & 1 \\ b & c & 1 \\ c & a & 1 \end{vmatrix} = 0.$$

(Used in solving 130.)

I. Solution by the PROPOSER.

First. *The given conditions are necessary*, for if two triangles abc, xyz , be similar, then $\frac{a-b}{x-y} = \frac{b-c}{y-z} = \frac{c-a}{z-x} = r \dots (1)$, for this expresses not only that the homologous sides are proportional, but also that the homologous angles are equal. (1) is equivalent to the equations

$$\begin{aligned} a-b &= r(x-y) \\ b-c &= r(y-z) \quad \dots (2). \\ c-a &= r(z-x) \end{aligned}$$

Multiplying the equations (2) by x, y , and z , respectively, and adding, we get

$$\Sigma(a-b)z = r \Sigma(x-y)z = 0 \dots (3),$$

or $\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ x & y & z \end{vmatrix} = 0 \dots (4)$, which is therefore necessary at least, if the triangles are similar.